

INVESTIGATION OF SELECTIVE BINDING OF MIANSERIN ENANTIOMERS WITH CYCLODEXTRINS

Nino Jajanidze¹, Ketevan Museridze¹, Bezhan Chankvetadze¹

*¹Chair of Physical and Analytical Chemistry, Department of Chemistry, School of Exact and Natural Sciences,
Iv. Javakishvili Tbilisi State University, Tbilisi 0179, Georgia
Email: nino.jajanidze392@ens.tsu.ge*

A significant part of biological processes is based on chiral recognition. Therefore, these recognition mechanisms are widely studied. One of the valuable instrumental methods for such studies is capillary electrophoresis (CE). Since neither electrophoretic nor electroosmotic mobility alone is enantioselective migration for enantiomers, addition of a chiral selector is necessary to separate them. For this purpose, native and substituted cyclodextrins (CD) can be used. CDs can bind enantiomers selectively and thus lead to a difference in their migration speed. Due to the high efficiency of CE, even the binding constant ratio as low as 1.01 is sufficient for the baseline separation of enantiomers. This cannot be achieved with any other separation methods [1].

The aim of this research was to study the affinity pattern of enantiomers of an antidepressant drug Mianserin with various, native and substituted CDs. α -, β - and γ - CDs were used as native CDs. These CDs consist of 6, 7, and 8 glucopyranose rings respectively. Therefore, their cavity sizes are different. The enantiomers were baseline separated using β - and γ -CDs, however, no separation was observed for α -CD. To study the migration order, the enantiomers were fractionated using high-performance liquid chromatography (HPLC). Then CE analysis was performed using the sample enriched with an enantiomer.

Various other substituted CDs were used, derivatives of all 3 native Cyclodextrins:
Hexakis(2,3-di-O-methyl-6-sulfo)-alpha-cyclodextrin sodium salt. (*HDMSACD*)
Heptakis(2,3-di-O-methyl-6-sulfo)-beta-cyclodextrin sodium salt. (*HDMSBCD*)
Octakis(2,3-di-O-methyl-6-sulfo)-gamma-cyclodextrin sodium salt. (*ODMSGCD*)
Heptakis(2,3-di-O-methyl-beta-cyclodextrin sodium salt.
2-Monocarboxymethylated beta-cyclodextrin
Hydroxypropylated-gamma-cyclodextrin (HP- γ -CD)
Peracetylated-beta-cyclodextrin (PA- β -CD)
The migration order is the same for all the above CDs.

References:

[1] P. Peluso, B.; Chankvetadze *Electrophoresis* **2021**, *42*, 1676-1708.